

Mathematical Journal of Okayama University

Volume 47, Issue 1

2005

Article 18

JANUARY 2005

Some Metric Invariants of Spheres and Alexandrov Spaces II

Nobuyuki Sochi*

*Okayama University

Copyright ©2005 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

Some Metric Invariants of Spheres and Alexandrov Spaces II

Nobuyuki Sochi

Abstract

A metric invariant a_k is defined, and we have that $a_k(X) \leq a_k(S^n)$ holds in an Alexandrov space X with curvature ≥ 1 ([So]). And the borderline case when $a_{2p-1}(X) = a_{2p-1}(S^n)$ and $a_k(S^n)$ are studied.

KEYWORDS: Metric Invariants, Alexandrov Spaces, Spheres, Borderline Cases

Math. J. Okayama Univ. **47** (2005), 193–202

SOME METRIC INVARIANTS OF SPHERES AND ALEXANDROV SPACES II

NOBUYUKI SOCHI

ABSTRACT. A metric invariant a_k is defined, and we have that $a_k(X) \leq a_k(S^n)$ holds in an Alexandrov space X with curvature ≥ 1 ([So]). And the borderline case when $a_{2p-1}(X) = a_{2p-1}(S^n)$ and $a_k(S^n)$ are studied.

1. INTRODUCTION

In the previous paper ([So]) we introduced a metric invariant $a_k(X)$ for a compact metric space X , and studied the explicit value of $a_k(S^n)$ for an n -dimensional round sphere S^n with radius 1. Furthermore, we studied the behavior of $a_k(X)$ for an Alexandrov space X with curvature ≥ 1 . In the present paper we continue to study the above invariant a_k and give answers to some problems that are conjectured in [So]. We begin with recalling the definition of a_k and results obtained in [So]. The distance between $x, y \in X$ will be denoted by $\text{dist}(x, y)$.

Definition 1.1. *For a positive integer k , we define the metric invariant $a_k(X)$ of X as follows:*

$$(1.1) \quad a_k(X) = \min_{x_1, \dots, x_k \in X} \max_{x \in X} \frac{1}{k} \sum_{i=1}^k \text{dist}(x, x_i).$$

In the previous paper we were concerned with $a_k(S^n)$ and got the explicit value of $a_k(S^1)$. A k -tuple (x_1, \dots, x_k) of points $x_i (i = 1, \dots, k)$ of S^1 located in counterclockwise order is called a configuration, where each x_i is called a vertex of the configuration.

Theorem 1.1. (1) *For $k = 2p - 1$, we have*

$$(1.2) \quad a_k(S^1) = \frac{2p^2 - 2p + 1}{(2p - 1)^2} \pi.$$

$a_k(S^1)$ is realized if and only if a configuration (x_1, \dots, x_k) of k points is equally spaced in S^1 , and $\max_{x \in S^1} (1/k) \sum_{i=1}^k \text{dist}(x, x_i)$ is attained exactly at the antipodal points of $x_i (1 \leq i \leq k)$.

(2) *For $k = 2p$, we have*

$$(1.3) \quad a_k(S^1) = \frac{1}{2} \pi.$$

$a_k(S^1)$ is realized if and only if (x_1, \dots, x_{2p}) consists of pairs of antipodal points, and in the case we have $(1/k) \sum_{i=1}^k \text{dist}(x, x_i) \equiv \pi/2$.

Now in this paper we complete the following theorem for general dimension n .

Theorem 1.2. (1) For $k = 2p - 1$, we have

$$(1.4) \quad a_k(S^n) = a_k(S^1) = \frac{2p^2 - 2p + 1}{(2p - 1)^2} \pi.$$

$a_{2p-1}(S^n)$ is realized if and only if $\{x_1, \dots, x_{2p-1}\}$ is located on a great circle S^1 and gives an equally spaced configuration after rearranging the order of $\{x_1, \dots, x_{2p-1}\}$, and $\max_{x \in S^n} (1/k) \sum_{i=1}^k \text{dist}(x, x_i)$ is attained exactly at the antipodal points of $x_i (1 \leq i \leq k)$.

(2) For $k = 2p$, we have

$$(1.5) \quad a_k(S^n) = \frac{1}{2} \pi.$$

Moreover, $a_k(S^n)$ is realized if and only if the set $\{x_1, \dots, x_{2p}\}$ consists of pairs of antipodal points, and in the case we have $(1/k) \sum_{i=1}^k \text{dist}(x, x_i) \equiv \pi/2$.

Theorem 1.2(2) is already proved in section 4 of [So]. Theorem 1.2(1) for $k = 3$ is also proved in section 3 of [So]. We give a proof of Theorem 1.2(1) for all $k = 2p - 1$ in section 2 of this paper. In our proof another metric invariant k -extent $xt_k(X)$ introduced by Grove and Markvorsen plays an important role. Recall that for an integer $k \geq 2$, $xt_k(X)$ is defined as follows:

$$(1.6) \quad xt_k(X) = \max_{x_1, \dots, x_k \in X} \binom{k}{2}^{-1} \sum_{i < j} \text{dist}(x_i, x_j).$$

Now we give some results of [G-M] on $xt_k(S^n)$ which are necessary for our proof.

Theorem 1.3 (Grove-Markvorsen). For all $n \geq 1$ and $k \geq 2$ we have

$$(1.7) \quad xt_k(S^n) = xt_k(S^1) = \pi / \left(2 - \left[\frac{k+1}{2} \right]^{-1} \right).$$

Those points that realize $xt_k(S^n)$ all lie on a great circle except for antipodal pairs.

More specifically, $xt_k(S^1)$ is given as follows:

$$(1.8) \quad xt_{2p}(S^1) = \frac{p}{2p-1} \pi,$$

$$(1.9) \quad xt_{2p-1}(S^1) = \frac{p}{2p-1}\pi.$$

$xt_{2p}(S^1)$ is realized if and only if $\{x_1, \dots, x_{2p}\}$ consists of pairs of antipodal points.

Now we explain the separation condition as follows. A configuration (x_1, \dots, x_k) in S^1 which does not contain any pair of antipodal points is said to satisfy the separation condition if the following is satisfied: For any x_i the line through x_i and the origin separates $(x_1, \dots, x_k) \setminus \{x_i\}$ into sets of equal cardinality. Note that k is odd in this case. Then $xt_{2p-1}(S^1)$ is realized if and only if $\{x_1, \dots, x_{2p-1}\}$ consists of pairs of antipodal points together with a configuration of points satisfying the separation condition.

For $k = 2p - 1$, $a_k(S^n)$ is related to $xt_k(S^n)$ by an inequality

$$(1.10) \quad a_k(S^n) \geq \pi - 2k^{-2} \binom{k}{2} xt_k(S^n) = \frac{2p^2 - 2p + 1}{(2p-1)^2} \pi,$$

as is shown in section 2.

Our proof of Theorem 1.2 depends also on a theorem of K. Kiyohara in [K] : For an equally spaced configuration (x_1, \dots, x_n) of a great circle S^1 of S^2 , a function $f_{x_1, \dots, x_k}(x) = \sum_{i=1}^k \text{dist}(x, x_i)$ on S^2 attains its maximum when and only when $x = \bar{x}_l (l = 1, \dots, k)$.

Let X be an Alexandrov space with curvature ≥ 1 and we want to compare $a_k(X)$ with $a_k(S^n)$. In the previous paper we got the following theorem by using the generalized Toponogov comparison theorem([Pe]).

Theorem 1.4. *Let X be an n -dimensional Alexandrov space with curvature ≥ 1 , then we have*

$$(1.11) \quad a_k(X) \leq a_k(S^n).$$

Especially, we have

$$(1.12) \quad a_{2p}(X) \leq a_{2p}(S^n) = \frac{\pi}{2}.$$

In the case where equality holds in (1.11) for $k = 3$ we showed that X is isometric to a double spherical suspension, where the generalized Toponogov comparison theorem played an important role(see section 5 of [So]). In the present paper we improve Theorem 1.4 and show the following theorem.

Theorem 1.5. *Let X be an n -dimensional Alexandrov space with curvature ≥ 1 . Suppose $a_{2p-1}(X) = a_{2p-1}(S^n)$. Then X is isometric to the unit sphere S^n .*

2. PROOF OF THEOREM 1.2(1)

In this section we give a proof of Theorem 1.2(1). First we will show for $k = 2p - 1$

$$(2.1) \quad a_k(S^n) \geq \frac{2p^2 - 2p + 1}{(2p - 1)^2} \pi.$$

In the proof we apply a result of Grove-Markvorsen on k -extent $xt_k(S^n)$ of the sphere (see Theorem 1.3). Recall that for $k = 2p - 1$ the k -extent $xt_k(S^n) = 1/\binom{k}{2} \max_{y_1, \dots, y_k \in S^n} \sum_{i < j} \text{dist}(y_i, y_j)$ of the sphere is equal to $p(p - 1)\pi / \binom{2p-1}{2} = p\pi / (2p - 1)$. Let x_1, \dots, x_k be points on S^n that realize $a_k(S^n)$, i.e., $a_k(S^n) = \max_{x \in S^n} 1/k \sum_{i=1}^k \text{dist}(x, x_i)$. We denote by \bar{x} the antipodal point of $x \in S^n$. Then for any \bar{x}_j ($1 \leq j \leq k$) we have

$$(2.2) \quad ka_k(S^n) \geq \sum_{i=1}^k \text{dist}(x_i, \bar{x}_j).$$

Adding (2.2) with respect to j we obtain

$$\begin{aligned} (2.3) \quad k^2 a_k(S^n) &\geq \sum_{j=1}^k \sum_{i=1}^k \text{dist}(x_i, \bar{x}_j) \\ &= k^2 \pi - 2 \sum_{i < j} \text{dist}(x_i, x_j) \\ &\geq k^2 \pi - 2 \max_{y_1, \dots, y_k \in S^n} \sum_{i < j} \text{dist}(y_i, y_j) \\ &= k^2 \pi - 2 \binom{k}{2} xt_k(S^n) = (2p^2 - 2p + 1)\pi. \end{aligned}$$

Hence we have (2.1). To complete the proof of Theorem 1.2(1) we have to show the following assertions:

- (1) $a_k(S^n) = (2p^2 - 2p + 1)\pi / (2p - 1)^2$.
- (2) If $\{x_1, \dots, x_k\}$ realizes $a_k(S^n)$, then $\{x_1, \dots, x_k\}$ is located on a great circle S^1 and a configuration obtained by rearranging $\{x_1, \dots, x_k\}$ is equally spaced on S^1 .
- (3) For an equally spaced configuration (x_1, \dots, x_k) , $f_{x_1, \dots, x_k}(x) = \sum_{i=1}^k \text{dist}(x, x_i)$, $x \in S^n$ takes the maximum exactly at $x = \bar{x}_j$ for all j .

To show these assertions we need the following theorem.

Theorem 2.1. *Let (x_1, \dots, x_k) be an equally spaced configuration on a great circle S^1 in S^n . The function $f(x) = \sum_{i=1}^k \text{dist}(x, x_i)$ takes its maximum at $x \in S^n$ when and only when $x = \bar{x}_l$ ($1 \leq l \leq k = 2p - 1$).*

Indeed, making use of Theorem 2.1 we may show (1), (2), and (3) as follows. First consider a configuration (x_1, \dots, x_k) as in Theorem 2.1, then we have $\max_{x \in S^n} f_{x_1, \dots, x_k} = \sum_{i=1}^k \text{dist}(x_i, \bar{x}_l) = (2p^2 - 2p + 1)\pi / (2p - 1)$. It follows that $a_k(S^n) \leq (2p^2 - 2p + 1)\pi / (2p - 1)^2$ and the assertion (1) holds. Next we show the assertion (2). Suppose $\{x_1, \dots, x_k\}$ realizes $a_k(S^n)$, i.e., $\max_{x \in S^n} f_{x_1, \dots, x_k} = (2p^2 - 2p + 1)\pi / (2p - 1)$. Then equality holds in the above inequalities (2.3), $\{x_1, \dots, x_k\}$ also realizes $xt_k(S^n)$, i.e., they are lying on a great circle S^1 except for some antipodal pairs by Theorem 1.3. Note, however, that $a_{2p-1}(S^n)$ cannot be realized if $\{x_1, \dots, x_k\}$ contains an antipodal pair. Indeed, suppose that $\{x_1, \dots, x_{2m-1}\} (1 \leq m < p)$ is on S^1 and $\{x_{2m}, \dots, x_{2p-1}\}$ consists of $(p - m)$ antipodal pairs. Then we have

$$\begin{aligned} \max_{x \in S^n} f_{x_1, \dots, x_{2p-1}}(x) &\geq \max_{x \in S^1} f_{x_1, \dots, x_{2p-1}}(x) \\ &\geq (p - m)\pi + (2m - 1)a_{2m-1}(S^1) \\ &= (p - m)\pi + \frac{2m^2 - 2m + 1}{2m - 1}\pi \\ &= \frac{(2p - 1)m + 1 - p}{2m - 1}\pi \\ &> \frac{2p^2 - 2p + 1}{2p - 1}\pi. \end{aligned}$$

It follows that $\{x_1, \dots, x_k\}$ is located on a great circle S^1 . We may also assume that (x_1, \dots, x_k) is a configuration in S^1 . Then $f_{x_1, \dots, x_k}(x)$ attains the maximum $ka_k(S^1) = \frac{2p^2 - 2p + 1}{2p - 1}\pi$ at $x = \bar{x}_j \in S^1$ by (2.3). By restricting f_{x_1, \dots, x_k} to S^1 we see that (x_1, \dots, x_k) is equally spaced in S^1 by Theorem 1.1. Then assertion (3) also follows from Theorem 2.1.

Now we give some remarks about a proof of Theorem 2.1. It suffices to consider the case $n = 2$ to prove Theorem 2.1. By virtue of Theorem 1.1 we need only to show that $f_{x_1, \dots, x_k}(x)$ cannot take a maximum at a point $x \in S^2 \setminus S^1$. For $k = 3$ we gave a proof of the theorem by showing that $f_{x_1, \dots, x_k}(x)$ admits no critical points in $S^2 \setminus S^1$ (see section 3 of [So]). But for $k = 2p - 1 > 3$ the behavior of critical points of $f_{x_1, \dots, x_k}(x)$ is rather complicated and it is not so clear whether the above approach works for general $k = 2p - 1$. Then K. Kiyohara gave a simple and ingenious proof of theorem 2.1 which will be presented in the appendix.

3. PROOF OF THEOREM 1.5

Let X be an n -dimensional Alexandrov space with curvature ≥ 1 . Recall that we have an inequality

$$(3.1) \quad a_{2p-1}(X) \leq a_{2p-1}(S^n) = \frac{2p^2 - 2p + 1}{(2p - 1)^2} \pi$$

by Theorem 1.2 and Theorem 1.4. In this section we show that X is isometric to the round sphere S^n of radius 1 when equality holds in (3.1). First we recall the notion of the spherical suspension and the spherical join ([B-G-P]).

Definition 3.1. *The spherical suspension of a metric space Y is the quotient space*

$$(3.2) \quad \Sigma_1 Y = Y \times [0, \pi] / \sim,$$

where the equivalence relation \sim is given by $(x_1, a_1) \sim (x_2, a_2) \Leftrightarrow x_1 = x_2, 0 < a_1 = a_2 < \pi$ or $a_1 = a_2 = 0$ or $a_1 = a_2 = \pi$, and is equipped with the canonical metric

$$(3.3) \quad \cos \operatorname{dist}(\hat{x}_1, \hat{x}_2) = \cos a_1 \cos a_2 + \sin a_1 \sin a_2 \cos \operatorname{dist}(x_1, x_2),$$

where we set $\hat{x}_1 = (x_1, a_1)$, $\hat{x}_2 = (x_2, a_2)$.

Definition 3.2. *The spherical join of X and Y is defined as*

$$(3.4) \quad X * Y = X \times Y \times [0, \pi/2] / \sim,$$

where $(x_1, y_1, a_1) \sim (x_2, y_2, a_2) \Leftrightarrow x_1 = x_2, y_1 = y_2, 0 < a_1 = a_2 < \pi/2$ or $a_1 = a_2 = 0$, $x_1 = x_2$ or $a_1 = a_2 = \pi/2$, $y_1 = y_2$, and is equipped with the canonical metric

$$(3.5) \quad \begin{aligned} & \cos \operatorname{dist}((x_1, y_1, a_1), (x_2, y_2, a_2)) \\ &= \cos a_1 \cos a_2 \cos \operatorname{dist}(x_1, x_2) + \sin a_1 \sin a_2 \cos \operatorname{dist}(y_1, y_2). \end{aligned}$$

Further, we define $\Sigma_k Y = \Sigma_{k-1}(\Sigma_1 Y)$ to be the k -times repeated spherical suspension. Then for an Alexandrov space X with curvature ≥ 1 we have $X = \Sigma_k Y$ for some Alexandrov space Y with curvature ≥ 1 if and only if S^{k-1} is isometrically embedded in X ([G-W]). Hence the k -times repeated spherical suspension $\Sigma_k Y$ is isometric to the spherical join $S^{k-1} * Y$.

In the previous paper, in the case of $k = 3$ we showed that X is isometric to $\Sigma_2 Z$ for some Alexandrov space Z with curvature ≥ 1 . First we show that X is isometric to the spherical suspension $\Sigma_1 Y$ in the same manner as in the case of $k = 3$ for completeness.

Lemma 3.1. *Let X be an n -dimensional Alexandrov space with curvature ≥ 1 . Suppose $a_{2p-1}(X) = a_{2p-1}(S^n) = \frac{2p^2 - 2p + 1}{(2p - 1)^2} \pi$. Then X is isometric to the spherical suspension $\Sigma_1 Y$, where Y is an $(n - 1)$ -dimensional Alexandrov space with curvature ≥ 1 .*

Proof. By the maximal diameter theorem([G-P2]), it suffices to show that $\text{diam}X$ is equal to π . Let $\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_{2p-1}$ be points on S^n that realize $a_{2p-1}(S^n)$. We may assume that $(\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_{2p-1})$ is an equally spaced configuration in a great circle S^1 . Take a point $\tilde{p} \in S^n$ different from the antipodals of $\tilde{x}_i (i = 1, 2, \dots, 2p-1)$. Take a regular point $p \in X$. We denote by S_p the space of directions of X at p . Then $\Sigma_1 S_p$ is isometric to S^n , and we identify $\Sigma_1 S_p$ (resp. S_p) with $S^n = \Sigma_1 S_{\tilde{p}}$ (resp. $S_{\tilde{p}} = S^{n-1}$). Let $x_i \in X$ be a point $\exp_p \bar{c}_{v_i}(\text{dist}(\tilde{p}, \tilde{x}_i)) = c_{v_i}(\text{dist}(\tilde{p}, \tilde{x}_i))$, where \bar{c}_{v_i} is a minimal geodesic in S^n emanating from \tilde{p} to \tilde{x}_i with initial direction $v_i \in S_{\tilde{p}} = S_p = S^{n-1} (i = 1, 2, \dots, 2p-1)$ and c_{v_i} is a quasigeodesic in X emanating from p with initial direction v_i (see section 5 of [So]). Take a point $x_0 \in X$ such that

$$\begin{aligned} a_{2p-1}(x_1, x_2, \dots, x_{2p-1}) &:= \max_{x \in X} \frac{1}{2p-1} \sum_{i=1}^{2p-1} \text{dist}(x, x_i) \\ &= \frac{1}{2p-1} \sum_{i=1}^{2p-1} \text{dist}(x_0, x_i). \end{aligned}$$

Let $\gamma_0 : [0, \text{dist}(p, x_0)] \rightarrow X$ be a minimal geodesic from p to x_0 , and set $\tilde{x}_0 = \exp_{\tilde{p}}^{S^n}(\text{dist}(p, x_0)\dot{\gamma}_0(0))$. Then by the generalized Toponogov comparison theorem for $\triangle p x_i x_0$ and $\triangle \tilde{p} \tilde{x}_i \tilde{x}_0$ ([Pe], see also [So]), we have

$$\text{dist}(x_0, x_i) \leq \text{dist}(\tilde{x}_0, \tilde{x}_i) \quad (i = 1, 2, \dots, 2p-1).$$

Then we have

$$\begin{aligned} (3.6) \quad a_{2p-1}(X) &\leq a_{2p-1}(x_1, x_2, \dots, x_{2p-1}) = \frac{1}{2p-1} \sum_{i=1}^{2p-1} \text{dist}(x_0, x_i) \\ &\leq \frac{1}{2p-1} \sum_{i=1}^{2p-1} \text{dist}(\tilde{x}_0, \tilde{x}_i) \leq a_{2p-1}(\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_{2p-1}) \\ &= a_{2p-1}(S^n) = a_{2p-1}(X). \end{aligned}$$

It follows that

$$(3.7) \quad a_{2p-1}(X) = a_{2p-1}(S^n) = \frac{1}{2p-1} \sum_{i=1}^{2p-1} \text{dist}(\tilde{x}_0, \tilde{x}_i),$$

and we obtain for any i

$$(3.8) \quad \text{dist}(x_0, x_i) = \text{dist}(\tilde{x}_0, \tilde{x}_i) \quad (i = 1, 2, \dots, 2p-1).$$

Then from Theorem 1.1 \tilde{x}_0 is the antipodal point of some \tilde{x}_i , namely, we have $\text{dist}(\tilde{x}_0, \tilde{x}_i) = \pi$, and hence $\text{dist}(x_0, x_i) = \pi$ for some $x_i (1 \leq i \leq 2p-1)$. \square

The following lemma is given in the previous paper ([So]).

Lemma 3.2. *Suppose $X = \Sigma_1 Y$, where Y is an $(n-1)$ -dimensional Alexandrov space with curvature ≥ 1 and $\text{diam} Y < \pi$ and $n \geq 2$. Let $x_1, x_2 \in X$ be the pole points of the spherical suspension $X = \Sigma_1 Y$. Then there is no pair of points whose distance is π except for x_1, x_2 .*

Next we show that X is isometric to $\Sigma_2 Z$ if $\dim X \geq 2$ as in the case of $k = 3$. By Lemma 3.2 we have the following lemma.

Lemma 3.3. *Let X be an n -dimensional Alexandrov space with curvature ≥ 1 and $n \geq 2$. Suppose $a_{2p-1}(X) = a_{2p-1}(S^n) = \frac{2p^2-2p+1}{(2p-1)^2}\pi$. Then X is isometric to $\Sigma_2 Z$, where Z is an $(n-2)$ -dimensional Alexandrov space with curvature ≥ 1 .*

Proof. By Lemma 3.1 we may write $X = \Sigma_1 Y$. Suppose $\text{diam} Y < \pi$. In the proof of Lemma 3.1 we may take a point p as an arbitrary regular point of X , and the set of regular points is dense in X . If the base point $p \in X$ is shifted, the points $x_1, x_2, \dots, x_{2p-1}$ that realize $a_{2p-1}(X)$ can be moved. Then there exists another pair of points $x_0, x_i (i = 1, 2, \dots, 2p-1)$ whose distance is equal to π . This contradicts Lemma 3.2. Therefore, we have $\text{diam} Y = \pi$ and $X = \Sigma_2 Z$. \square

We prepare one more lemma.

Lemma 3.4. *Let X be an n -dimensional Alexandrov space with curvature ≥ 1 . Suppose $a_{2p-1}(X) = a_{2p-1}(S^n) = \frac{2p^2-2p+1}{(2p-1)^2}\pi$. If X is isometric to $S^{k-1} * Y^{n-k} (1 \leq k \leq n-1)$, where Y^{n-k} is an $(n-k)$ -dimensional Alexandrov space with curvature ≥ 1 , then $\text{diam} Y^{n-k} = \pi$.*

Proof. Suppose $\text{diam} Y^{n-k} < \pi$. Take any points $(x, y, t_1), (x', y', t_2) \in X (x, x' \in S^{k-1}, y, y' \in Y^{n-k}, 0 \leq t_1, t_2 \leq \pi/2)$, and set

$$l = \text{dist}((x, y, t_1), (x', y', t_2)).$$

Then we will show that $l = \pi$ holds exactly when $t_1 = t_2 = 0$ and $\text{dist}(x, x') = \pi$, namely, $(x, y, t_1), (x', y', t_2)$ are antipodal pair of S^{k-1} . Indeed, by the distance formula (3.5) we have

$$\begin{aligned} -1 &= \cos l \\ (3.9) \quad &= \cos t_1 \cos t_2 \cos \text{dist}(x, x') + \sin t_1 \sin t_2 \cos \text{dist}(y, y') \\ &\geq \cos(\pi + t_1 - t_2) \geq -1. \end{aligned}$$

Then we have $t_1 = t_2 = 0$ because of $\cos(\text{dist}(y, y')) > -1$, and also $\text{dist}(x, x') = \pi$ holds.

Now in the proof of Lemma 3.1 we may choose a point p arbitrarily as long as p is regular. Since $\text{rad} X = a_1(X) \geq a_{2p-1}(X) = (2p^2 - 2p + 1)\pi/(2p - 1)^2 > \pi/2$, X is homeomorphic to S^n by the radius sphere theorem, and

the set S of non regular points is a closed set of dimension $\leq n - 2$ ([G-P1]). Also we may choose identification between S_p and $S_{\bar{p}} = S^{n-1}$ up to isometries of S^{n-1} . If the point $p \in X$ is shifted or $v_1, v_2, \dots, v_{2p-1}$ are rotated around p in $S_p = S^{n-1}$, the points $x_1, x_2, \dots, x_{2p-1}$ that realize $a_{2p-1}(X)$ can be moved outside of S^{k-1} . Then there exists another pair of points $x_0, x_i (i = 1, 2, \dots, 2p-1)$ with $\text{dist}(x_0, x_i) = \pi$ such that they are not antipodal pair in S^{k-1} . This contradicts $\text{diam} Y^{n-k} < \pi$. It follows that $\text{diam} Y^{n-k} = \pi$ holds. \square

Now we show that X is isometric to S^n .

Proof of Theorem 1.5. By Lemma 3.1 and Lemma 3.3 we have $X = S^0 * Y^{n-1} = S^1 * Y^{n-2}$. Next we assume that $X = S^{k-1} * Y^{n-k}$ holds for $k (1 \leq k \leq n-1)$. By Lemma 3.4 we have $\text{diam} Y^{n-k} = \pi$. It follows that $X = S^k * Y^{n-k-1}$. By induction on k we see that X is isometric to $S^{n-2} * Y^1$. Since $\text{rad} X = \text{rad}(\Sigma_1 Y^{n-1}) = \text{rad} Y^{n-1}$, we have $\text{rad} Y^{n-1} > \pi/2$ (see [G-P1]). It follows that $\text{rad} Y^1 > \pi/2$ and Y^1 is homeomorphic to the circle S^1 . By Lemma 3.4 $\text{diam} Y^1 = \pi$ and therefore Y^1 is isometric to S^1 . It follows that X is isometric to S^n . This completes the proof of Theorem 1.5. \square

Remark 3.1. If $a_{2p-1}(X)$ is close to $a_{2p-1}(S^n) = (2p^2 - 2p + 1)\pi / (2p - 1)^2 > \pi/2$, then X is homeomorphic to S^n since $\text{rad} X > \pi/2$.

ACKNOWLEDGMENT

I would like to express my deepest gratitude to professors, Takashi Sakai, Atsushi Katsuda, and Kazuyoshi Kiyohara for the encouragement and helpful suggestions, as well as for teaching me all the necessary background.

REFERENCES

- [B-G-P] Y. Burago-M. Gromov-G. Perelman. *Alexandrov spaces with curvature bounded below I*. Russ. Math. Surveys. **47**, 1-58(1992).
- [G-M] K. Grove-S. Markvorsen. *New extremal problems for the Riemannian recognition program via Alexandrov geometry*. J. Amer. Math. **8**, 1-28(1995).
- [G-P1] K. Grove-P. Petersen. *A radius sphere theorem*. Invent. Math. **112**, 577-583(1993).
- [G-P2] K. Grove-P. Petersen. *On the excess of metric spaces and manifolds*. Preprint.
- [G-W] K. Grove- F. Wilhelm. *Hard and soft packing theorems*. Ann. of Math. **142**, 213-237(1995).
- [K] K. Kiyohara. *Appendix to "Some metric invariants of spheres and Alexandrov spaces II"*. to appear in Math. J. Okayama Univ. **47**.
- [P] G. Perelman. *Alexandrov spaces with curvature bounded below II*. Preprint.
- [Pe] A. Petrunin. *Quasigeodesics in multidimensional Alexandrov spaces*. Diploma, University of Illinois(1995).
- [P-P] G. Perelman- A. Petrunin. *Extremal subsets in Alexandrov spaces and the generalized Lieberman theorem*. St. Petersburg Math J. **5**, 215-227(1994).

- [S] S. Shteingold. *Covering Radii and Paving Diameters of Alexandrov Spaces*. J. Geom. Anal. **8**, 613-627(1998).
- [So] N. Sochi. *Some metric invariants of spheres and Alexandrov spaces I*. Math. J. Okayama Univ.**46**, 163-182(2004).

NOBUYUKI SOCHI

THE GRADUATE SCHOOL OF NATURAL SCIENCE AND TECHNOLOGY

OKAYAMA UNIVERSITY

OKAYAMA 700-8530, JAPAN

e-mail address: iputiko@yahoo.co.jp

(Received September 29, 2004)

(Revised December 28, 2004)